

Journal of Psychopharmacology

<http://jop.sagepub.com/>

Cannabinoids in bipolar affective disorder: a review and discussion of their therapeutic potential

C. H. Ashton, P. B. Moore, P. Gallagher and A. H. Young

J Psychopharmacol 2005 19: 293

DOI: 10.1177/0269881105051541

The online version of this article can be found at:

<http://jop.sagepub.com/content/19/3/293>

Published by:

 SAGE

<http://www.sagepublications.com>

On behalf of:



[British Association for Psychopharmacology](#)

Additional services and information for *Journal of Psychopharmacology* can be found at:

Email Alerts: <http://jop.sagepub.com/cgi/alerts>

Subscriptions: <http://jop.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

Citations: <http://jop.sagepub.com/content/19/3/293.refs.html>

>> [Version of Record](#) - May 11, 2005

[What is This?](#)

Cannabinoids in bipolar affective disorder: a review and discussion of their therapeutic potential

C. H. Ashton *Department of Psychiatry, University of Newcastle upon Tyne, Royal Victoria Infirmary, Newcastle upon Tyne, UK.*

P. B. Moore *Department of Psychiatry, University of Newcastle upon Tyne, Royal Victoria Infirmary, Newcastle upon Tyne, UK.*

P. Gallagher *Department of Psychiatry, University of Newcastle upon Tyne, Royal Victoria Infirmary, Newcastle upon Tyne, UK.*

A. H. Young *Department of Psychiatry, University of Newcastle upon Tyne, Royal Victoria Infirmary, Newcastle upon Tyne, UK.*

Abstract

Bipolar affective disorder is often poorly controlled by prescribed drugs. Cannabis use is common in patients with this disorder and anecdotal reports suggest that some patients take it to alleviate symptoms of both mania and depression. We undertook a literature review of cannabis use by patients with bipolar disorder and of the neuropharmacological properties of cannabinoids suggesting possible therapeutic effects in this condition. No systematic studies of cannabinoids in bipolar disorder were found to exist, although some patients claim that cannabis relieves symptoms of mania and/or depression. The cannabinoids Δ^9 -tetrahydrocannabinol (THC) and cannabidiol (CBD) may exert sedative, hypnotic, anxiolytic, antidepressant, antipsychotic and

anticonvulsant effects. Pure synthetic cannabinoids, such as dronabinol and nabilone and specific plant extracts containing THC, CBD, or a mixture of the two in known concentrations, are available and can be delivered sublingually. Controlled trials of these cannabinoids as adjunctive medication in bipolar disorder are now indicated.

Keywords

bipolar disorder, cannabidiol, cannabinoids, cannabis, CBD, depression, dronabinol, mania, nabilone, tetrahydrocannabinol, THC

Introduction

The treatment of bipolar affective disorder (BAD) remains problematic despite several guidelines or consensus statements (Sachs *et al.*, 2000; Geddes and Goodwin, 2001; Goodwin, 2003; Lloyd *et al.*, 2003). The mean time to relapse after the first episode is 5 years (Geddes *et al.*, 2003) and periods of remission shorten as the illness progresses, regardless of treatment. Most patients with BAD are prescribed a combination of drugs, all of which have their disadvantages. Lithium, although efficacious, has limited effectiveness because of low acceptance and occurrences of mania on withdrawal. Many anticonvulsants can produce unacceptable side-effects (Porter *et al.*, 1999; Ashton and Young, 2003). Sodium valproate, the most commonly prescribed mood stabilizer, carries risks in women of childbearing age (Committee on Safety of Medicines, 2003; Goodwin and Sachs, 2004). Lamotrigine,

although effective in bipolar depression, requires careful dosage control to prevent skin complications, which may prove to be serious. Conventional antidepressants and electroconvulsive therapy can induce mood elevation, which may progress to rapid mood cycling. Antipsychotic drugs have many undesirable effects and the atypical antipsychotics quetiapine, olanzapine and risperidone have all been reported to induce mania in some cases (Mishra *et al.*, 2004). Psychosocial measures have been shown to complement medication, but they remain at an early stage of development and their widespread use is limited by available resources.

Thus, there is a clear need to explore new ways of managing bipolar disorder. Patient reports and observations, backed by known pharmacology, suggest that the cannabis derivatives Δ^9 -tetrahydrocannabinol (THC) and cannabidiol (CBD) may have mood stabilizing properties. The present study aimed to review the evidence for this. The use of controlled substances in medicine is

Corresponding author: Professor C. H. Ashton, Department of Psychiatry, University of Newcastle upon Tyne, Royal Victoria Infirmary, Newcastle upon Tyne NE1 4LP, UK.
Email: c.h.ashton@ncl.ac.uk

widespread, especially in children with psychological difficulties and in pain management. Nevertheless, the consequences of extending the use of controlled substances need careful consideration.

It is well known that there is a high prevalence of comorbid drug abuse in people with BAD (Brown *et al.*, 2001). A 61% lifetime prevalence of substance abuse in Bipolar I patients and 48% in Bipolar II patients has been reported compared to 6% in the general population (Regier *et al.*, 1990). Some studies have provided data on individual drugs that are abused by these patients (Estroff *et al.*, 1985; Miller *et al.*, 1989; Regier *et al.*, 1990; Marken *et al.*, 1992; Mueser *et al.*, 1992; Sonne *et al.*, 1994; Winokur *et al.*, 1998). The results indicate high rates of lifetime use of cannabis (30–64%) and stimulants (amphetamines 31–39%, cocaine 15–39%) and lower rates for opiates (6–25%). The extent to which bipolar patients use cannabis as self-medication is not clear, although anecdotal reports suggest that some patients find it alleviates both depression (Gruber *et al.*, 1996) and mania (Grinspoon and Bakalar, 1998). Although cannabis can cause adverse effects, including psychosis and mania, some cannabinoids have properties that could be of value in psychiatric disorders, and a literature review was therefore undertaken to investigate their therapeutic potential in bipolar affective disorder.

Methods

Electronic searches for relevant papers were performed, employing Medline (1966 to present), Embase (1980 to present), ISI Web of Science (1990 to present) and Psychoinfo (earliest available to present). Search terms were 'bipolar', 'manic depression', 'mania', 'antidepressant', 'antimanic', 'mood stabilizer', 'cannabinoid', 'tetrahydrocannabinol', 'THC', 'cannabidiol', 'CBD', 'cannabis', 'marijuana', 'nabilone' and 'dronabinol'.

In addition, Medline reviews and investigations of pharmacological, psychiatric and therapeutic effects of cannabis/cannabinoids (1970–2003) were consulted and a manual searching of all relevant articles was performed.

Results

The literature search revealed no systematic studies of the therapeutic use of cannabis or cannabinoids in BAD, although there are several anecdotal reports. Grinspoon and Bakalar (1998) described five cases in which cannabis appeared to alleviate mania. For example, one woman with BAD quoted in their report chose cannabis over alcohol to control her manic behaviour: 'A few puffs of this herb and I can be calm ... this drug seems harmless compared to other drugs I have tried, including tranquillisers and lithium'. A husband, describing his wife with BAD said: 'My wife functions much better when she uses marijuana. When she is hypomanic, it relaxes her, helps her sleep, and slows her speech down. When she is depressed and would otherwise lie in bed all day, the marijuana makes her more active ... Lithium is also effective, but it doesn't always keep her in control'.

Personal observation of a patient attending the local outpatients also indicated an apparent antimanic effect of cannabis. The patient was a 39-year-old male who had been diagnosed 10 years previously as having BAD. His illness mainly took the form of manic episodes for which he had a history of five hospital admissions. These episodes were difficult to control because the patient was intolerant of antipsychotic drugs, including quetiapine and risperidone, and non-compliant with lithium and sodium valproate. Diazepam controlled his symptoms but he often used up his 2-week prescription for 30 mg daily in 1 week.

A recent manic episode was associated with a severe behaviour disturbance involving a further possible detention order. The psychiatrist was called for a home visit, which he made some hours later. To his surprise, he found the patient calm, almost serene, sitting tranquilly in an armchair smoking a cannabis 'spliff'. (He offered the psychiatrist one of the same, which was declined). It was clear that the cannabis was responsible for the rapid change in the patient's behaviour. He maintained that, over the years, he had taken mainly cannabis, sometimes moderate amounts of alcohol, occasionally 'street' benzodiazepines, and infrequently heroin to regulate his mood.

Gruber *et al.* (1996) described five cases in which marijuana appeared to produce a direct antidepressant effect. Three of these patients had BAD and all but one found that marijuana relieved their depression better than standard antidepressant drugs. Two surveys of medicinal cannabis use in California, where this use is legalized, showed that 15–27% of patients were prescribed it for mood disorders, including depression, post-traumatic stress disorder, BAD and attention deficit disorder resistant to conventional pharmacotherapy (Gieringer, 2003).

It is noteworthy that, in the anecdotal reports, cannabis was not taken for the 'high' sought by recreational users and it is possible that its effects are different when taken in subeuphoric doses for medical reasons, such as in multiple sclerosis or pain conditions (Randall, 1991; Hodges, 1993). The effects are most probably due to cannabinoids present in cannabis smoke, including Δ^9 -THC, CBD and possibly others, which have been less studied. Patients' accounts and the advances in the understanding of cannabinoid physiology suggest that they may have a therapeutic potential in BAD (Pertwee, 1999a,b).

Pharmacological basis of cannabinoid effects: the endocannabinoid system

THC and cannabinoid CB_1 receptors THC is the major psychoactive agent present in cannabis, and its primary metabolite, 11-OH-THC, is even more potent (Maykut, 1985; McPartland and Russo, 2001). These cannabinoids are agonists of endogenous cannabinoid CB_1 receptors that are present in the brain, spinal cord and peripheral nerves. CB_1 receptors are widely distributed throughout the brain (Table 1) and are present in the cerebral cortex, including the cingulate cortex, hippocampus, basal amygdala, corpus striatum and other areas possibly involved in the pathophysiology of BAD and its emotional and cognitive components (Drevets *et al.*, 1997; Strakowski *et al.*, 1999; Altshuler *et al.*, 2000; Phillips *et al.*,

Table 1 Localization of cannabinoid CB₁ receptors

Density	Localization
Very dense	Basal ganglia
	Cerebellum
	Hippocampus
Dense	Cerebral cortex ^a
	Hippocampus
	Corpus striatum
Moderate	Hypothalamus ^a
	Basal amygdala ^a
	Central grey substance
	Nucleus of solitary tract
	Spinal cord
	Peripheral nerve terminals
Sparse	Thalamus
	Pons and Medulla
	Some non-neural tissues, including spleen and testes

^aReceptor density in the cingulate cortex, hypothalamus and amygdala is relatively greater in the human brain than in the same areas of rat and monkey brain (Herkenham, 1995; Pertwee, 1997).

2003; Surguladze *et al.*, 2003). CB₁ receptors belong to a family of G-protein coupled receptors that includes receptors for aminergic neurotransmitters (noradrenaline, dopamine, serotonin and acetylcholine) and act through second messenger systems. CB₂ receptors are similar to CB₁ receptors but are present mainly in immune cells in the periphery and are not considered further here.

Activation of the CB₁ receptor (Fig. 1) inhibits adenylate cyclase and decreases the production of cAMP (3,5-adenosine monophosphate) (Pertwee, 1997), an action which affects many intracellular processes and ultimately affects intracellular neurotransmission (Shiloh *et al.*, 1999). CB₁ receptors also modulate transneuronal ion channels. They are negatively coupled to calcium channels (N and P/Q type) and inhibit the inward flow of calcium ions, decreasing the release of neurotransmitters, either excitatory or inhibitory, at presynaptic nerve terminals (Pertwee, 1997). At the same time, CB₁ activation enhances the outward flow of potassium ions (through A-type potassium channels), a G-protein coupled event that may also depend on inhibition of cAMP production (Deadwyler *et al.*, 1995). The result is inhibition of neuronal depolarization, decreased action potential generation and hence reduced impulse propagation.

CBD and anandamides The endogenous ligands for cannabinoid receptors, both CB₁ receptors in the nervous system and CB₂ receptors in peripheral tissues, are a family of arachidonic acid derivatives, sometimes termed endocannabinoids (Pertwee, 1999a,b). The two that appear to be of most physiological importance are arachidonyl ethanolamide (anandamide) and 2-arachidonyl glycerol (2-AG). Anandamide is present in the brain in the same areas as CB₁ receptors. It is enzymatically synthesized in cell membranes, binds to CB₁ receptors (Van der Stelt and Di Marzo,




Figure 1 Schematic diagram of signal transduction mechanisms stimulated by CB₁ receptors. The CB₁ receptor (1) is coupled to a second messenger G_{i/o} protein. Via this protein, activation of the receptor inhibits the enzyme adenylate cyclase (2) and decreases the production of cAMP (3). Via the G-protein, the inward flow of calcium ions is blocked (4), decreasing release of neurotransmitters (5). Also via the G-protein, the outward flow of potassium ions is enhanced (6), resulting in decreased neuronal firing and decreased impulse transmission (7). Stimulation of the G-protein also activates MAP kinase (8), affecting intracellular gene expression. Other receptors on the same neurone (for monoamines and/or opioids) may activate their own G-proteins but share a common adenylate cyclase, which they may stimulate (9) or inhibit (10). Anandamide is released in the post-synaptic membrane and acts retrogradely as an agonist on presynaptic CB₁ receptors (Howlett, 1995; Pertwee, 1997; Ameri, 1999; Joy *et al.*, 1999; Van der Stelt and di Marzo, 2003; Alger, 2004).

2003) and, in animal models, shows many of the actions of THC (Stein *et al.*, 1996; Martin and Cone, 1999). However, unlike THC, the effects of anandamide are short-lived, lasting less than 15 min after intravenous injection in the rat (Stein *et al.*, 1996) because it is rapidly inactivated by enzymatic hydrolysis and removed from its site of action by neuronal uptake mechanisms (Joy *et al.*, 1999; Pertwee, 1997, 1999b; Piomelli *et al.*, 2000; Alger, 2004). In addition, anandamide is synthesized and released at discrete loci on demand by neural activity or depolarization of postsynaptic membranes and then acts retrogradely as an agonist on presynaptic CB₁ receptors (Piomelli *et al.*, 2000; Christie and Vaughan, 2001; Wilson and Nicol, 2001; Van der Stelt and Di Marzo, 2003; Alger, 2004). By contrast, the exogenous cannabinoid THC is widely distributed, reaching all areas of CB₁ receptors, is very slowly eliminated (Agurell *et al.*, 1986) and produces effects lasting several hours (Maykutt, 1985).

CBD binds only minimally to CB₁ receptors and is usually described as non-psychoactive. However, the clinical observations described below suggest that it has antipsychotic, anxiolytic, anti-convulsant and other psychological effects (Zuardi *et al.*, 1995; Mechoulam *et al.*, 2002). Its mode of action is not fully understood but CBD has recently been shown to block the reuptake of

anandamide (Bisogno *et al.*, 2001) and to inhibit its enzymatic hydrolysis (Mechoulam *et al.*, 2002). CBD also reduces the hydroxylation of THC to its more psychoactive metabolite, 11-OH-THC (McPartland and Russo, 2001). It has been shown to inhibit serotonin reuptake and to increase catecholamine activity in rat brain synaptosomes (McPartland and Russo, 2001), an action also shown by anandamide (Steffens and Feuerstein, 2004). In addition, CBD is a potent antioxidant agent and is protective against glutamate toxicity, an action which is not affected by cannabinoid receptor antagonists (Mechoulam *et al.*, 2002). The possible contribution of each of these actions to the psychological effects of CBD is not clear.

The discovery of endocannabinoids and the realization that these are the biological ligands of cannabinoid receptors has opened a whole new vista in cannabinoid pharmacology. A system of cannabinoid receptors and endocannabinoids appears to modulate many important physiological processes (Di Marzo *et al.*, 1998). These processes have yet to be clearly defined but evidence is already accumulating that endocannabinoids are involved in the modulation of brain reward systems (Gardner, 1999), mood, anxiety and sleep (Musty *et al.*, 1995), pain (Pertwee, 2001), cognition and memory (Terranova *et al.*, 1995, 1996), appetite (Williams and Kukham, 1999; Di Marzo *et al.*, 2001), endocrine activity (Mendelson and Mello, 1999), cardiovascular regulation (Randall and Kendall, 1998) and other vital functions (Musty *et al.*, 1995; Ameri, 1999). The basic function of the endogenous system appears to be the regulation of interneuronal signalling, involving complex interactions with many neurotransmitters and neuromodulators, including monoamines, acetylcholine, opioids, GABA and glutamate (Ameri, 1999).

Psychological effects of THC

The psychological effects of cannabis and THC have been described by many authors (Paton and Pertwee, 1973; Ashton, 1999a; Johns, 2001). It is important to note that many of these are biphasic and bidirectional, depending on dose, mode of administration, environment, expectation, personality, degree of tolerance and other individual factors, as well as time-frame (Paton and Pertwee, 1973; Ashton *et al.*, 1981; Ashton, 1999b). Thus, acute effects in normal subjects can include euphoria or dysphoria, relaxation or anxiety, excitation followed by sedation, heightened perception followed by perceptual distortion, and increased motor activity followed by incoordination. Synthetic THC (dronabinol) and nabilone, a synthetic cannabinoid related to THC, exert similar actions depending on dosage and the other factors mentioned above. In healthy subjects under placebo-controlled laboratory conditions, THC (5 mg and 10 mg smoked in herbal cigarettes) was shown to produce relaxation with decreased subjective ratings of anxiety, tension and depression (Ashton *et al.*, 1981). However, D'Souza *et al.*, 2004) recently found that intravenous infusions of THC (2.5 mg and 5 mg) produced mild and transient schizophrenia-like symptoms, anxiety, detachment, perceptual distortion and cognitive impairment.

Patients using cannabis or synthetic THC compounds in moderate doses for chronic pain conditions or multiple sclerosis have

reported improvement of mood and increased general well-being and mental health, as well as alleviation of their other symptoms (Martyn *et al.*, 1995; Notcutt *et al.*, 1997; Ashton, 1999b; Williams and Evans, 2000; Wade *et al.*, 2003; Svendson *et al.*, 2004). A few controlled studies have shown anxiolytic effects of nabilone in some patients (Glass *et al.*, 1980; Fabre and McLendon, 1981; Ilaria *et al.*, 1981) and an antidepressant effect of THC in cancer patients (Regelson *et al.*, 1976; Russo *et al.*, 2003).

Many of the adverse effects of cannabis (usually attributed to its THC content) result from relatively high dose or chronic use. Cannabis can cause an acute psychosis in previously normal individuals, but those with mental illness are more vulnerable (Johns, 2001). Such reactions are dose-related and appear to be becoming more common with the present-day recreational use of potent cannabis varieties such as 'skunk' and netherweed (Wylie *et al.*, 1995). Heavy cannabis use can also lead to an acute functional psychosis with marked hypomanic features (Rottenburg *et al.*, 1982; Johns, 2001). In patients with BAD, the duration of cannabis use is associated positively with the duration of manic, but not depressive, episodes (Strakowski *et al.*, 2000) and substance abuse in general appears to increase the severity of the illness (Cassidy *et al.*, 2001) and to increase suicide rate (Dalton *et al.*, 2003).

Cannabis is a well-known risk factor for schizophrenia and may precipitate the illness in genetically predisposed individuals (Johns, 2001). It aggravates positive symptoms in schizophrenia and may antagonize the effects of antipsychotic drugs (Negrete and Gill, 1999). A large number of studies, as reviewed by Arsenault *et al.* (2004) and Macleod *et al.* (2004), have implicated a dose-related association between the use of cannabis in childhood and adolescence with later development in young adulthood of schizophrenia, depression, violence and antisocial behaviour, use of other illicit drugs, lower educational attainment, and psychological distress. Whether or not these associations are causal are debated by the above authors.

Psychological effects of CBD

There is some evidence that CBD, which constitutes up to 40% of cannabis extracts, has anxiolytic, hypnotic, antipsychotic and anticonvulsant actions (Zuardi and Guimaraes, 1997; Mechoulam *et al.*, 2002). CBD antagonizes the anxiety, intoxication liability and psychotic-like symptoms produced by high doses of THC in normal subjects (Zuardi *et al.*, 1982; Russo, 2003) and has similar anxiolytic effects to diazepam in a simulated public speaking test (Zuardi and Guimaraes, 1997). Anxiolytic effects have also been demonstrated in animal models, including the behaviour of rodents on the elevated plus maze (Guimaraes *et al.*, 1990). In this test, the action of CBD, administered alone, was dose-dependent and biphasic, similar to many other cannabinoid effects (Sulcova *et al.*, 1998). Biphasic hypnotic effects in rats have also been demonstrated (Monti, 1997) and CBD significantly increased sleeping time compared to placebo in insomniacs (Carlini and Cunha, 1981).

Antipsychotic effects of CBD were suggested by the observation that it acted in a similar way to haloperidol in animal tests predictive of antipsychotic activity (Zuardi *et al.*, 1991, 1995). A placebo-controlled case study of a patient with schizophrenia who was

intolerant of haloperidol showed antipsychotic effects of high-dose oral CBD with 60–69% improvement in scores on the Brief Psychiatric Rating Scale and Interactive Observation Scale for Psychiatric Inpatients after 4 weeks of CBD therapy (Zuardi *et al.*, 1995). Preliminary results with CBD in additional schizophrenic patients are reported as promising (Gerth *et al.*, 2002).

Anticonvulsant actions of CBD, comparable to those of diphenylhydantoin and other drugs that are clinically effective in major seizures, have been shown in a variety of animal models (Consroe and Snyder, 1986; Consroe and Sandyk, 1992). The effects are not reversed by CB₁ antagonists, indicating that they are not CB₁ receptor mediated. A small placebo-controlled clinical study of oral CBD as an add-on therapy in 15 patients with uncontrolled secondary generalized epilepsy with temporal focus was conducted by Cunha *et al.* (1980). Of the eight patients who received CBD over 4 months, four remained almost seizure-free and three others showed partial improvement, whereas the patients taking placebo showed no change.

Pharmacokinetic factors

When administered orally, the absorption of both THC and CBD is slow and erratic. Peak plasma concentrations are not reached for 2–6 h and the biological availability is 4–12% for THC (Grotenhermen, 2003) and 13–19% for CBD (Mechoulam *et al.*, 2002). Both cannabinoids undergo extensive first pass metabolism in the liver and THC is also degraded by stomach acids. By contrast, inhaled cannabinoids reach peak plasma concentrations within minutes and have a bioavailability of approximately 35% for both THC and CBD. For medicinal purposes, other modes of administration have been investigated and sublingual liquid solutions appear to be well absorbed, producing rapid effects comparable to inhalation (Whittle *et al.*, 2001; Grotenhermen, 2003; Wade *et al.*, 2003). Using a sublingual spray of THC and CBD, Wade *et al.* (2003) found that it was possible for subjects with pain conditions or multiple sclerosis to self-titrate small doses that relieved pain and muscle spasms without inducing intoxication.

After absorption, both THC and CBD are sequestered in fatty tissues from which they are only slowly released (the tissue half-life is 5–7 days). Both cannabinoids form a large number of metabolites, which are gradually eliminated over days or weeks in the urine and faeces (Gold, 1992). There may be complex interactions between the two cannabinoids. CBD inhibits some cytochrome P450 enzymes and may inhibit the conversion of THC to its active 11-hydroxy metabolite (McPartland and Russo, 2001), but Zuardi *et al.* (1982) found no effect on THC levels in humans when the two cannabinoids were administered together. By contrast, THC and its metabolites, and even CBD on repeated administration, increase cytochrome P450 activity through enzyme induction (Grotenhermen, 2003).

Discussion

Despite the sparse anecdotal data in humans and the absence of controlled clinical trials, the evidence discussed above shows that

Table 2 Comparison of some effects of Δ⁹-tetrahydrocannabinol (THC) and cannabidiol (CBD)

Actions	THC	CBD
Agonist action on CB ₁ receptors	+	–
Inhibition of anandamide reuptake and hydrolysis	–	+
Anxiolytic	+ ^a	+
Psychotropic	+	–
Antipsychotic	–	+ ^b
Anticonvulsant	–	+
Antidepressant	(+) ^c	–
Sedative/hypnotic	+	+
Antinociceptive	+	+
Neuroprotective (inhibition of glutamate release)	+	+
Antiemetic	+	–
Appetite stimulant	+	No data
Cardiovascular effects ^d	+	+

^aTHC is anxiolytic in some doses, but can be anxiogenic in higher doses or in drug-naïve individuals. ^bCBD also antagonizes some psychotropic effects of THC. ^cShown in one study in cancer patients (Regelson *et al.*, 1976). ^dTHC causes tachycardia and hypotension; CBD can cause bradycardia and hypotension.

both THC and CBD have pharmacological properties that could be therapeutic in patients with BAD. Furthermore, the available pharmacokinetic evidence indicates optimal methods of administration and dosage control. The underlying pathophysiology of BAD is unknown, but these cannabinoids, especially when used in combination, have several characteristics (Table 2) in common with drugs known to benefit this disorder, including antidepressants, antipsychotics, anticonvulsants (mood-stabilizers) and anxiolytics.

THC, in some conditions and doses, has anxiolytic, hypnotic and antidepressant effects with improvement in mood and general well-being in normal subjects, and in patients with pain conditions, multiple sclerosis or cancer (Regelson *et al.*, 1976; Glass *et al.*, 1980; Ashton *et al.*, 1981; Fabre and McLendon, 1981; Ilaria *et al.*, 1981; Paton and Pertwee, 1981; Martyn *et al.*, 1995; Notcutt *et al.*, 1997; Ashton, 1999b; Wade *et al.*, 2003). These actions could be helpful in BAD, especially in depressive phases, which are often accompanied by anxiety (Goodwin and Sachs, 2004). CBD antagonizes the psychotic-like effects and intoxication liability produced by high doses of THC and has anxiolytic, hypnotic and anticonvulsant actions of its own in addition to a protective effect against glutamate toxicity (Cunha *et al.*, 1980; Carlini and Cunha, 1981; Consroe and Snider, 1986; Guimaraes *et al.*, 1990; Consroe and Sandyk, 1992; Zuardi *et al.*, 1995; Zuardi and Guimaraes, 1997; Gerth *et al.*, 2002; Mechoulam *et al.*, 2002; Russo, 2003). These actions do not appear to be mediated by CB₁ receptors but may result from enhancement of the endogenous anandamide system and effects on THC metabolism (Mechoulam *et al.*, 2002; McPartland and Russo, 2001). As well as adding to the anxiolytic effects of THC, the antipsychotic effects of CBD could be therapeutic in bipolar patients with psychotic symptoms, and the anticonvulsant and protective effects against glutamate toxicity may have a mood-stabilizing action similar to some other anticonvulsants of proven value in BAD (Porter *et al.*, 1999; Ashton and

Young, 2003). In addition, both THC and CBD have extremely low toxicity (British Medical Association, 1997; Mechoulam *et al.*, 2002).

Cannabinoids have already been tested for therapeutic effects in acute and chronic pain conditions and multiple sclerosis (Wade *et al.*, 2003; Svendsen *et al.*, 2004). The evidence suggests that a placebo-controlled trial of cannabinoids as adjunctive therapy in BAD should now be undertaken. Such a trial might start with a pilot investigation in treatment-resistant bipolar patients who remain symptomatic despite standard medications, choosing patients over the age of 18 years who have used cannabis previously (but who undertake to abstain from cannabis during the trial). Standardized plant extracts containing THC and CBD in combination and matching placebo have been available for clinical research since 1988 (GW Pharmaceuticals plc, Salisbury, UK). These could be self-administered as a 1 : 1 THC : CBD mixture or placebo and delivered by metered dose pump action aerosol spray as described by Wade *et al.* (2003). These authors found that the product was well tolerated and that side-effects were minimal in patients with various neurological disorders. Bipolar patients could self-titrate their preferred dosage to control symptoms and dosage would be minimized by limiting the amount contained in each spray to 2.5 mg of cannabinoid and the total dosage in each daily container to 120 mg cannabinoids. Thus, the maximum amount of THC obtainable daily would be 60 mg: a single modern cannabis 'spliff' contains 60–150 mg THC or more (Ashton, 1999b). Treatment periods would possibly be for 4 weeks, perhaps in a crossover active treatment/placebo design. Assessments would include clinical ratings of mania and depression scores, subjective rating scales, neuropsychological performance and a record of adverse effects. The results would provide information on optimal dosage regimes, duration of treatment, adverse effects and other factors.

Possible adverse effects that would require close monitoring in such a trial include the precipitation of hypomania, mania and psychosis, although these effects are unlikely to be significant with small dose preparations and a 50% CBD content in the medication. Neurocognitive function, which is already impaired in BAD (El-Badri *et al.*, 2001; Ferrier and Thompson, 2002) may be further compromised by THC (Solowij, 1998). On the other hand, better symptom control with the THC/CBD preparation may improve cognition. Additive effects may occur with hypnotics, sedatives and alcohol. Induction of cytochrome P450 enzymes may result in interactions with drugs metabolized by the same enzymes, including many antidepressants and antipsychotics. However, these enzymes are already induced in BAD patients who smoke tobacco or take cannabis. Two patients who stopped or reduced tobacco and/or cannabis consumption when on clozapine or olanzapine experienced adverse effects due to increased plasma levels of the drugs, necessitating dosage adjustment (Zullino *et al.*, 2002). A possible interaction between lithium and marijuana was reported in one case resulting in elevated serum lithium levels, which dropped when the patient stopped using marijuana (Ratey *et al.*, 1981). The interaction was attributed to slowed gut motility caused by marijuana which increased lithium absorption.

Tolerance and dependence can result from chronic use of cannabis and withdrawal effects occur on ceasing use (Ashton, 1999a). However, little tolerance appears to develop to the putative

therapeutic effects that have been studied. Some patients have found nabilone still to be effective for pain relief after 2–3 years of regular use (Notcutt *et al.*, 1997) and patients taking plant-based cannabinoid extracts long-term for pain have not so far reported tolerance (Whittle *et al.*, 2001). Any withdrawal problems could be minimized by tapering dosage if use was no longer required. Similar to cannabis, THC has abuse potential and precautions may be needed to limit patients' overuse of the cannabinoid aerosols.

In conclusion, BAD is often poorly controlled by existing drugs and often involves a polypharmacological medley, including lithium, anticonvulsants, antidepressants, antipsychotics and benzodiazepines. Many patients take street drugs in addition, including cannabis, amphetamines, cocaine and illicitly obtained benzodiazepines in an attempt to control their symptoms. Some claim that such self-medication is superior to the drugs prescribed by psychiatrists. There are good pharmacological reasons for believing that the prescription of synthetic cannabinoids or standardized plant extracts may have a therapeutic potential in BAD. We suggest that the time is ripe for carefully managed trials of prescribed cannabinoids to determine whether they are of value as adjunctive drugs in bipolar patients whose symptoms are not adequately controlled by standard medications.

References

Agurell A, Halldin M, Lindgren J E, Ohlsson A, Widman M, Gillespie H, Hollister L (1986) Pharmacokinetics and metabolism of Δ^1 -tetrahydrocannabinol and other cannabinoids with emphasis on man. *Pharmacol Rev* 38: 21–43

Alger B E (2004) Endocannabinoids: getting the message across. *Proc Natl Acad Sci* 101: 8512–8513

Altshuler L L, Bartzokis G, Grieder T, Curran J, Jimenez T, Leight K, Wilkins J, Gerner R, Mintz J (2000) An MRI study of temporal lobe structures in men with bipolar disorder or schizophrenia. *Biol Psychiatry* 48: 147–162

Ameri A (1999) The effects of cannabinoids on the brain. *Prog Neurobiol* 58: 315–348

Arsenault L, Cannon M, Witton J, Murray R M (2004) Causal association between cannabis and psychosis: examination of the evidence. *Br J Psychiatry* 184: 110–117

Ashton H (1999a) Cannabis in palliative care. *CME Bull Palliat Med* 1: 73–77

Ashton H (1999b) Adverse effects of cannabis and cannabinoids. *Br J Anaesth* 83: 637–649

Ashton H, Young A H (2003) GABA-ergic drugs: exit stage left, enter stage right. *J Psychopharmacol* 17: 174–178

Ashton H, Golding J F, Marsh V R, Millman J E, Thompson J W (1981) The seed and the soil: effect of dosage, personality and starting state on the response to Δ^9 -tetrahydrocannabinol in man. *Br J Clin Pharmacol* 12: 705–720

Bisogno T, Hanus L, De Petrocellis L, Tchilibon S, Ponde D, Brandi I, Moriello A S, Davis J B, Mechoulam R, Di Marzo V (2001) Molecular targets for cannabidiol and its synthetic analogues: effect on vanilloid VR1 receptors and on the cellular uptake and enzymatic hydrolysis of anandamide. *Br J Pharmacol* 134: 845–852

British Medical Association (1997) *Therapeutic Uses of Cannabis*. Harwood Publishers, London

Brown E, Suppes T, Adinoff B, Thomas NR (2001) Drug abuse and bipolar disorder: comorbidity or misdiagnosis? *J Affect Dis* 65: 105–115

Carlini E A, Cunha J M (1981) Hypnotic and antiepileptic effects of cannabidiol. *J Clin Pharmacol* 21: 417S–427S

Cassidy F, Ahearn E P, Carroll B J (2001) Substance abuse in bipolar disorder. *Bipolar Disord* 3: 181–188

Christie M J, Vaughan C W (2001) Cannabinoids act backwards. *Nature* 410: 527–30

Committee on Safety of Medicines (2003) Current Problems in Pharmacovigilance 29: 6

Consroe P, Sandyk R (1992) Potential role of cannabinoids for therapy of neurological disorders. In Murphy L, Bartke A (eds), *Marijuana/Cannabinoids: Neurobiology and Neuropsychology*. CRC Press, Boca Raton, FL, pp. 459–524

Consroe P, Snyder S R (1986) Therapeutic potential of cannabinoids for therapy of neurological disorders. In Mechoulam R (ed.), *Cannabinoids as Therapeutic Agents*. CRC Press, Boca Raton, FL, pp. 21–49

Cunha J M, Carlini E A, Pereira A E, Ramos O L, Pimentel G, Gagliardi R, Sanvito E L, Lander N, Mechoulam R (1980) Chronic administration of CBD to healthy volunteers and epileptic patients. *Pharmacologia* 21: 175–185

Dalton E J, Cate-Carter T D, Mundo E, Parikh S V, Kennedy J L (2003) Suicide risk in bipolar patients: the role of co-morbid substance use disorders. *Bipolar Disord* 3: 58–61

Deadwyler S A, Hampson R E, Childers S R (1995) Functional significance of cannabinoid receptors in brain. In Pertwee R G (ed.), *Cannabinoid Receptors*. Academic Press Ltd, London, pp. 205–231

Di Marzo V, Goparaju S K, Wang L, Liu J, Batkai S, Jarai Z M, Fezza F, Miura G I, Palmiter R D, Sugiura T, Kunos G (2001) Leptin-regulated endocannabinoids are involved in maintaining food intake. *Nature* 410: 822–825

Di Marzo V, Melck D, Brisogno T, De Petrocellos L (1998) Endocannabinoids: endogenous cannabinoid receptor ligands with neuro-modulatory action. *Trends Neurosci* 21: 521–528

D'Souza D C, Perry E, MacDougall L, Ammerman Y, Cooper T, Wu, Y, Braley G, Gueorguieva R, Krystal J H (2004) The psychotomimetic effects of intravenous delta-9-tetrahydrocannabinol in healthy individuals: implications for psychosis. *Neuropsychopharmacology* 29: 1558–1572

Drevets W C, Price J L, Simpson J R (1997) Subgenual prefrontal cortex abnormalities in mood disorders. *Nature* 386: 824–827

El-Badri S M, Ashton C H, Moore P B, Marsh V R, Ferrier I N (2001) Electrophysiological and cognitive function in young euthymic patients with bipolar affective disorder. *Bipolar Disord* 3: 79–87

Estroff T W, Dackis C A, Golod M S, Pottash A L C (1985) Drug abuse and bipolar disorders. *Int J Psychiatry Med* 15: 37–40

Ferrier I N, Thompson J M (2002) Cognitive impairment in bipolar affective disorder: implications for the bipolar diathesis. *Br J Psychiatry* 180: 293–295

Fabre L F, McLendon D (1981) The efficacy and safety of nabilone (a synthetic cannabinoid) in the treatment of anxiety. *J Pharmacol Sci* 21: 377S–382S

Gardner E L (1999) Cannabinoid interaction with reward systems. In Nahas G G, Sutin K M, Harvey D J, Agurell S (eds), *Marijuana and Medicine*. Humana Press, Totowa, New Jersey, pp. 187–205

Geddes J R, Carney S M, Davies C (2003) Relapse prevention with antidepressant treatment in depressive disorders. *Lancet* 361: 643–651

Geddes J, Goodwin G (2001) Bipolar disorder: clinical uncertainty, evidence-based medicine and large-scale randomised trials. *Br J Psychiatry* 41 (Suppl): S191–S194

Gerth C W, Schulte-Lutter F, Mauss C, Ruhrmann S, Klosterkötter J, Leweke F M (2002) B113: the natural cannabinoid cannabidiol in the treatment of acute schizophrenia. *Schizophr Res* 53 (Suppl): 192

Gieringer D H (2003) The acceptance of medicinal marijuana in the US. *J Cannabis Ther* 3: 53–65

Glass R M, Uhlenhuth E H, Hartel F W, Schuster C R, Fischman M W (1980) A single dose study of nabilone, a synthetic cannabinoid. *Psychopharmacology* 71: 137–142

Gold M S (1992) Marijuana and hashish. In Winger G, Hoffman F G, Woods J H (eds). *A Handbook of Drug and Alcohol Abuse. The Biological Aspects*. Chapter 7. Oxford University Press, Oxford, pp. 117–131

Goodwin G M (2003) Evidence-based guidelines for treating bipolar disorder: recommendations from the British Association of Psychopharmacology. *J Psychopharmacol* 17: 149–173

Goodwin G, Sachs G (2004) Definitions: Diagnosis and Comorbidity. *Bipolar Disorder – Fast Facts*. Health Press, Oxford

Grinspoon L, Bakalar J B (1998) The use of cannabis as a mood stabilizer in bipolar disorder: anecdotal evidence and the need for clinical research. *J Psychoactive Drugs* 30: 171–177

Grotenhermen F (2003) Clinical pharmacokinetics. *J Cannabis Ther* 3: 3–51

Gruber A J, Pope H G, Brown M E (1996) Do patients use marijuana as an antidepressant? *Depression* 4: 77–80

Guimaraes F S, Chiaretti T M, Graeff F G, Zuardi A W (1990) Antianxiety effect of cannabidiol in the elevated plus-maze. *Psychopharmacology* 100: 558–559

Herkenham M (1995) Localisation of cannabinoid receptors in brain and periphery. In Pertwee R G (ed.), *Cannabinoid Receptors*. Academic Press, London, pp. 145–166

Hodges C (1993) I wish I could get it at the chemist's. *The Independent* 23rd February 1993.

Howlett A C (1995) Cannabinoid compounds and signal transduction mechanisms. In Pertwee R G (ed.), *Cannabinoid Receptors*. Academic Press, London, pp. 167–204

Ilaria R L, Thornby J I, Fann W E (1981) Nabilone, a cannabinol derivative, in the treatment of anxiety neurosis. *Curr Ther Res* 29: 943–949

Johns A (2001) The psychiatric effects of cannabis. *Br J Psychiatry* 178: 116–122

Joy J E, Watson S J, Benson J A (eds) (1999) *Marijuana and Medicine*. National Academy Press, Washington, DC

Lloyd A J, Harrison C L, Ferrier I N, Young A H (2003) The pharmacological treatment of bipolar affective disorder: practice is improving but could be better. *J Psychopharmacol* 17: 230–233

Macleod J, Oakes R, Copello A, Crome I, Egger M, Hickman M, Oppenkowski T, Stokes-Lampard H, Smith G D (2004) Psychological and social sequelae of cannabis and other illicit drug use by young people: a systematic review of longitudinal, general population studies. *Lancet* 363: 1579–1588

Marken P A, Stanislav S W, Lacombe S, Pierce C, Horstra R, Sommi R W (1992) Profile of a sample of subjects admitted to an acute care psychiatric facility with manic symptoms. *Psychopharmacol Bull* 28: 201–205

Martin B R, Cone E J (1999) Chemistry and pharmacology of cannabis. In Kalant H, Corrigan W A, Hall W, Smart R G (eds), *The Health Effects of Cannabis*. Centre for Addiction and Mental Health, Toronto, pp. 21–68

Martyn C N, Illis L S, Thom J (1995) Nabilone in the treatment of multiple sclerosis. *Lancet* 345: 579

Maykut M O (1985) Health consequences of acute and chronic marijuana use. *Prog Neuropsychopharmacol Biol Psychiatry* 9: 209–238

McPartland J M, Russo E B (2001) Cannabis and cannabis extracts: greater than the sum of their parts? *J Cannabis Ther* 1: 103–32

Mechoulam R, Parker L A, Gallily R (2002) Cannabidiol: an overview of some pharmacological aspects. *Clin Pharmacol* 42: 11S–19S

Mendelson J H, Mello N K (1999) Marijuana effects on pituitary and gonadal hormones. In Nahas G G, Sutin K M, Harvey D, Agurell S (eds), *Marijuana and Medicine*. Humana Press, Totowa, NJ, pp. 385–392

Miller F T, Busch F, Tanenbaum J H (1989) Drug abuse in schizophrenia and bipolar disorder. *Am J Drug Alcohol Abuse* 15: 291–295

Mishra A, Moore P B, Hobbs R (2004) Does quetiapine have mood altering properties? *J Psychopharmacol* 18: 281–284

Monti J M (1997) Hypnotic-like effects of cannabidiol in the rat. *Psychopharmacology* 55: 263–265

Mueser K T, Yarnold P R, Bellack A S (1992) Diagnostic and demographic correlates of substance abuse in schizophrenia and major affective disorder. *Acta Psychiatr Scand* 85: 105–114

Musty R E, Reggio P, Consroe P (1995) A review of recent advances in cannabinoid research and the 1994 international symposium on cannabis and the cannabinoids. *Life Sci* 56: 1933–1940

Negrete J C, Gill K (1999) Cannabis and schizophrenia: an overview of the evidence to day. In Nahas G G, Sutin K M, Harvey D, Agurell S (eds), *Marijuana and Medicine*. Humana Press, Totowa, NJ, pp. 671–682

Notcutt W, Price M, Chapman G (1997) Clinical experience with nabilone for chronic pain. *Pharmaceutical Sci* 3: 551–555

Paton W D M, Pertwee R G (1973) The actions of cannabis in man. In Nahas G G, Paton W D M (eds), *Marijuana: Chemistry, Pharmacology, Metabolism and Clinical Effects*. Pergamon Press, Oxford, pp. 735–38

Pertwee R G (1997) Pharmacology of cannabinoid CB₁ and CB₂ receptors. *Pharmacol Ther* 74: 129–180

Pertwee R G (1999a) Cannabinoid receptors and their ligands in brain and other tissues. In Nahas G G, Sutin K M, Harvey D, Agurell S (eds), *Marijuana and Medicine*. Humana Press, Totowa, NJ, pp. 177–185

Pertwee R G (1999b) Pharmacology of cannabinoid receptor ligands. *Curr Med Chem* 6: 635–664

Pertwee R G (2001) Cannabinoid receptors and pain. *Prog Neurobiol* 63: 569–611

Phillips M L, Drevets W C, Rauch S L, Lane R (2003) Neurobiology of emotion perception II: Implications for major psychiatric disorders. *Biol Psychiatry* 54: 515–528

Piomelli D, Giuffrida A, Calignano A, de Fonseca F R (2000) The endocannabinoid system as a target for therapeutic drugs. *Trends Pharm Sci* 21: 218–224

Porter R, Ferrier N, Ashton H (1999) Anticonvulsants as mood stabilisers. *Adv Psychiatr Treat* 5: 96–103

Randall R (ed.) (1991) *Muscle Spasm, Pain and Marijuana Therapy*. Galen Press, Washington, DC

Randall M D, Kendall D A (1998) Endocannabinoids: a new class of vasoactive substances. *Trends Pharm Sci* 19: 55–58

Ratey J J, Ciraulo D A, Shader R I (1981) Lithium and marijuana. *J Clin Psychopharmacol* 1: 32–33

Regelson W, Butler J R, Schulz J, Kirk T, Peek L, Green M L, Zalis M O (1976) Delta-9-THC as an effective antidepressant and appetite-stimulating agent in advanced cancer patients. In Braude M C, Szara S (eds), *The Pharmacology of Marijuana*. Raven Press, New York, NY, pp. 763–776

Regier D A, Farmer M E, Rae D S, Locke B Z, Keith S J, Judd L L, Goodwin F K (1990) Comorbidity of mental disorders with alcohol and other drug abuse. *J Am Med Assoc* 264: 2511–2518

Rottanburg D, Robins A H, Ben-Arie O, Teggin A, Elk R (1982) Cannabis-associated psychosis with hypomanic features. *Lancet* 2: 1364–1366

Russo E (2003) Cannabis: from pariah to prescription. *J Cannabis Ther* 3: 1–29

Sachs G S, Printz D J, Kahn D A (2000) The Expert Consensus Guideline Series: Medication treatment of bipolar disorder. *Postgrad Med Spec* No: 1–104

Shiloh R, Nutt D, Weizman A (1999) *Atlas of Psychiatric Chemotherapy*. Martin Dunitz, London

Solowij N (1998) *Cannabis and Cognitive Function*. Cambridge University Press, Cambridge

Sonne S C, Brady K T, Morton W A (1994) Substance abuse and bipolar affective disorder. *J Nerv Ment Dis* 182: 349–352

Steffens M, Feuerstein T J (2004) Receptor-independent depression of DA and 5-HT uptake by cannabinoids in rat neocortex – involvement of Na(+)/K(+)-ATPase. *Neurochem Int* 44: 529–538

Stein E A, Fuller S A, Edgemond W S, Campbell W B (1996) Physiological and behavioural effects of the endogenous cannabinoid, arachidonyl-ethanolamine (anandamide), in the rat. *Br J Pharmacol* 119: 107–114

Strakowski S M, DelBello M P, Sax K W (1999) Brain magnetic resonance imaging of structural abnormalities in bipolar disorder. *Arch Gen Psychiatry* 56: 254–260

Strakowski S M, DelBello M P, Fleck D E, Arndt S (2000) The impact of substance abuse on the course of bipolar disorder. *Biol Psychiatry* 48: 477–485

Sulcova E, Mechoulam R, Fride E (1998) Biphasic effects of anandamide. *Pharmacol Biochem Behav* 59: 347–353

Surguladze S, Keedwell P, Phillips M (2003) Neural systems underlying affective disorders. *Adv Psychiatr Treat* 9: 446–455

Svendsen K B, Jensen T S, Bach F W (2004) Does the cannabinoid dronabinol reduce central pain in multiple sclerosis? Randomised double blind placebo controlled crossover trial. *Br Med J* 329: 253–258

Terranova J P, Michaud J D, Le Fur G, Soubrie P (1995) Inhibiting long-term potentiation in rat hippocampal slice by anandamide and WIN55, 212-2: Reversal by SR1411716A, a selective antagonist of CB₁ cannabinoid receptors. *Naunyn-Schmiedeberg's Arch Pharmacol* 352: 576–579

Terranova J P, Storme J J, Lafon N, Perio A, Rinaldi-Ca J, Le Fur G, Soubrie P (1996) Improvement in memory in rodents by selective CB₁ cannabinoid receptor antagonist, SR141716 A. *Psychopharmacology* 126: 165–172

Van der Stelt M, Di Marzo V (2003) The endocannabinoid system in the basal ganglia and in the mesolimbic reward system: implications for neurological and psychiatric disorders. *Eur J Pharmacol* 480: 133–150

Wade D, Robson P, House H, Makela P, Aram J (2003) A preliminary controlled study to determine whether whole-plant cannabis extracts can improve intractable neurogenic symptoms. *Clin Rehabil* 17: 18–26

Whittle B A, Guy G W, Robson P (2001) Prospects for new cannabis-based prescription medicines. *J Cannabis Ther* 1: 183–205

Williams C M, Kukham T C (1999) Anandamide induces overeating: mediation by central cannabinoid (CB₁) receptors. *Psychopharmacology* 143: 315–317

Williamson E M, Evans F J (2000) Cannabinoids in clinical practice. *Drugs* 60: 1303–1314

Wilson R I, Nicol R A (2001) Endogenous cannabinoids mediate retrograde signalling at hippocampal synapses. *Nature* 410: 558–559

Winokur G, Turvey C, Skiskal H, Coryell W, Solomon D, Leon A, Mueller T, Endicott J, Maser J, Keller M (1998) Alcoholism and drug abuse in three groups – bipolar I, unipolars and their acquaintances. *J Affect Disord* 50: 81–89

Wylie A S, Scott R T A, Burnett S J (1995) Psychosis due to 'skunk'. *Br Med J* 311: 125

Zuardi A W, Guimaraes F S (1997) Cannabidiol as an anxiolytic and antipsychotic. In Mathre M L (ed.), *Cannabis in Medical Practice*. McFarland & Company, Jefferson, NC, pp. 133–141

Zuardi A W, Shirakawa I, Finkelfarb E, Karmol I G (1982) Action of cannabidiol on the anxiety and other effects produced by Δ⁹-THC in normal subjects. *Psychopharmacology* 76: 245–259

Zuardi A W, Rodrigues J A, Cunha J M (1991) Effects of cannabidiol in animal models predictive of antipsychotic activity. *Psychopharmacology* 104: 260–264

Zuardi A W, Morais S L, Guimaraes F S, Mechoulam R (1995) Antipsychotic effect of cannabidiol. *J Clin Psychiatry* 56: 485–486

Zullino D F, Delessert D, Eap C B, Preisij M, Baumann P (2002) Tobacco and cannabis smoking cessation can lead to intoxication with clozapine or olanzapine. *Int Clin Psychopharmacol* 17: 141–143